

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

city lot seven or eight American Goldfinches, but a search failed to disclose a single dead bird of any other species, though Yellow Warblers and English Sparrows at least are as abundant in the vicinity as the Goldfinches. I could obtain no information as to difference in the shelter sought by the different species which would account for the partiality exhibited by the storm.

JUNIUS HENDERSON, Boulder, Colo.

Golden Eagle (Aquila chrysaetos) at Cadiz, Ohio.—Mr. J. Bingham Bargar, who lives six miles south of Cadiz, wrote me recently in regard to a Golden Eagle he once shot and later had mounted. He writes: "It was first seen after a storm about December 1, 1887. We then saw it almost every day for a month. It lived on wild game, and when game was scarce it would take chickens. It finally killed a fine chicken, and I made an effort to trap it alive, but it broke the trap and got away. I followed it and was able to shoot it. It weighed fourteen pounds and twelve ounces, and measured seven feet, five and three-quarter inches from tip to tip of wings."

Concerning Passenger Pigeon (Ectopistes migratorius).—In a conversation once with Dr. Beal, of Scio College, who is the author of the Beal Law, he told me that a guide he met while on a vacation trip in northern Michigan in 1903, could possibly throw some light on the mysterious disappearance of the Passenger Pigeon. The guide claims that after a great storm a number of years ago, countless numbers of Wild Pigeons were thrown upon the shore of the lake. Delbert Burdett, a farmer, living near Cadiz, claims that while at work cutting timber near St. Clairsville, Belmont County, Ohio, in September, 1898, a flock of "thousands of Wild Pigeons" suddenly appeared, and in alighting covered a field several acres in extent. A number of old farmers saw the flock and all agreed that the birds were Wild Pigeons. Mr. Burdett is familiar with a large number of birds and I questioned him rather closely as to the Pigeons, and have reason to believe his story is correct.

Cadiz, Ohio.

HARRY B. McCONNELL.

Notes from Cadiz, Ohio.—Holboell's Grebe (Colymbus holbællii).—On February 6, 1905, some boys saw a strange bird fly against the telephone wires and fall to the ground in an alley in Cadiz, and as it was unable to continue its flight they picked it up and brought it to me to identify. It proved to be a Holbæll's Grebe, the first and only record I have of this bird for the county. The boys thought they might assist it to continue its journey by taking it up to the top of the Court House and pitching it out into space, expecting to see it take wing and fly away. This they did, but the